Table 20. Reported cases of HIV infection (not AIDS) for male adults and adolescents, by transmission category and race/ethnicity, cumulative through 2004—42 areas with confidential name-based HIV infection reporting

Transmission category	White, not Hispanic				Black, not Hispanic				Hispanic			
	2004		Cumulative through 2004 <sup>a</sup>		2004		Cumulative through 2004 <sup>a</sup>		2004		Cumulative through 2004 <sup>a</sup>	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact Injection drug use Male-to-male sexual contact and	5,729 506	68 6	41,771 4,660	66 7	3,459 1,004	37 11	22,485 10,417	34 16	2,212 1,144	44 23	10,399 4,709	43 19
injection drug use Hemophilia/coagulation disorder Heterosexual contact	453 17 246	5 0 3	4,314 341 2,094	7 1 3	240 9 1,173	3 0 13	2,871 103 9,089	4 0 14	171 1 562	3 0 11	948 24 2,195	4 0 9
Sex with injection drug user Sex with person with hemophilia Sex with HIV-infected transfusion	51 1	1 0	493 7	1 0	150 1	2 0	1,435 15	2 0	87 4	2 0	369 7	2 0
recipient Sex with HIV-infected person, risk factor not specified	5 189	0 2	31 1,563	0 2	8 1,014	0 11	77 7,562	0 11	4 467	<i>0</i> 9	14 1,805	<i>0 7</i>
Receipt of blood transfusion, blood components, or tissue Other/risk factor not reported	11	0	194	0	27	0	213	0	8	0	49	0
or identified  Total	1,477 <b>8,439</b>	18 <b>100</b>	9,644 <b>63,018</b>	15 <b>100</b>	3,460 <b>9,372</b>	37 <b>100</b>	21,528 <b>66,706</b>	32 <b>100</b>	978 <b>5,076</b>	19 <b>100</b>	5,895 <b>24,219</b>	24 <b>100</b>

	Asian/Pacific Islander				American Indian/Alaska Native				9	Total			
	2004		Cumulative through 2004 <sup>a</sup>		2004		Cumulative through 2004 <sup>a</sup>		2004b		Cumulative through 2004 <sup>a</sup>		
Transmission category	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%	
Male-to-male sexual contact	123	60	554	57	56	59	417	55	11,601	50	76,003	48	
Injection drug use	8	4	45	5	7	7	83	11	2,675	12	19,989	13	
Male-to-male sexual contact and													
injection drug use	6	3	21	2	5	5	100	13	878	4	8,298	5	
Hemophilia/coagulation disorder	1	0	2	0	0	0	0	0	29	0	474	0	
Heterosexual contact	16	8	66	7	8	8	49	6	2,014	9	13,564	9	
Sex with injection drug user	1	0	8	1	3	3	15	2	293	1	2,330	1	
Sex with person with hemophilia Sex with HIV-infected transfusion	0	0	0	0	0	0	0	0	6	0	29	0	
recipient Sex with HIV-infected person,	0	0	2	0	0	0	0	0	17	0	124	0	
risk factor not specified	15	7	56	6	5	5	34	4	1,698	7	11,081	7	
Receipt of blood transfusion,													
blood components, or tissue Other/risk factor not reported	0	0	3	0	0	0	2	0	46	0	468	0	
or identified	51	25	284	29	19	20	110	14	6,015	26	38,251	24	
Total	205	100	975	100	95	100	761	100	23,258	100	157,047	100	

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2004, the following 42 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Georgia, Idaho, Indiana, Iowa, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Puerto Rico, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

<sup>&</sup>lt;sup>a</sup> Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through December 2004. Cumulative total includes 1,368 males of unknown race or multiple races.

<sup>&</sup>lt;sup>b</sup> Total includes 71 males of unknown race or multiple races.